

# PUR TABS™

EFFERVESCENT SANITIZING / DISINFECTION TABLETS



## ACTIVE INGREDIENT:

Sodium dichloro-s-triazinetrione .....	48.21%*
Other Ingredients:.....	51.79%
Total .....	100.00%

\*Equivalent to 31.10% active chlorine by tablet weight. Refer to dilution chart for Available Chlorine concentrations

## KEEP OUT OF REACH OF CHILDREN

# DANGER

### See product container label for additional precautionary statements and first aid and full directions for use.

For use in Cleaning and Disinfection on hard non-porous surfaces in beverage and food processing plants, schools, hospitals, nursing homes, child care centers, restaurants, stores, veterinary clinics, zoos and aquariums, dairy farms, farms, poultry premises, industrial facilities, kennels, boarding facilities, laboratories, lab animal facilities, institutions, catering, kitchens, Intensive Care Unit, operating rooms, dental facilities, gyms, health clubs, and restrooms. Effective against *Clostridium difficile* spores. Effective against Hepatitis A virus, Hepatitis B virus and Hepatitis C virus.

**PUR TABS** is effective against the following micro-organisms on pre-cleaned, hard, non-porous, inanimate surfaces: *Salmonella enterica*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Staphylococcus epidermidis*, *Escherichia coli* O157:H7, *Staphylococcus aureus* - methicillin-resistant (MRSA) & glycopeptide-resistant (GRSA), carbapenem resistant *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Streptococcus pneumoniae*, vancomycin resistant *Enterococcus faecalis*, Poliovirus type 1, Herpes simplex virus type 1, Hepatitis A virus, Hepatitis B virus, Hepatitis C virus, Human Immunodeficiency Virus Type 1 (associated with AIDS), Influenza virus H1N1, respiratory syncytial virus, Canine Parvovirus, Newcastle Disease Virus, Pseudorabies, Canine Distemper Virus, Feline Calicivirus, Norovirus, Coxsackievirus, *Trichophyton interdigitale*, *Aspergillus fumigatus*, *Candida albicans*, *Mycobacterium bovis* (TB) and *Clostridium difficile* spores. Refer to Usage Table for solution concentration and contact times.

**PUR TABS** is designed to provide effective cleaning, and disinfection in areas where it is of prime importance in controlling the hazard of cross contamination on treated pre-cleaned, hard, non-porous, inanimate surfaces.

**PUR TABS** is a disinfectant that disinfects pre-cleaned, hard, non-porous, inanimate surfaces. This cleaning process may be accomplished with any cleaner solution including PUR TABS.

**PUR TABS** provides effective cleaning strength that will not dull high gloss floor finishes with repeated use.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**DANGER: Corrosive. Causes irreversible eye damage. Harmful if swallowed, inhaled, or absorbed through skin.** Do not get in eyes, on skin, or clothing. Avoid breathing dust. Wear chemical-resistant gloves and safety glasses or face shield when making up solution. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

### FIRST AID:

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **IF SWALLOWED:** Call a poison control center, or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

### IN THE EVENT OF A MEDICAL EMERGENCY CALL YOUR POISON CONTROL CENTER AT 1-800-222-1222.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

**PHYSICAL OR CHEMICAL HAZARDS: STRONG OXIDIZING AGENT:** Use only clean dry utensils. Mix only into water. Contamination with moisture, dirt, organic matter or other chemicals or any other foreign matter may start a chemical reaction with generation of heat, liberation of hazardous gases and possible generation of fire and explosion. Avoid any contact with flaming or burning material such as a lighted cigarette. Do not use this product in any chlorinating device which has been used with any inorganic or unstabilized chlorinating compounds (e.g., calcium hypochlorite). Such use may cause fire or explosion.

## DIRECTIONS FOR USE

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label and use strictly in accordance with precautionary statements and directions.**

**PUR TABS** is a Hospital Use Disinfectant. As a Healthcare disinfectant it is effective against standard Gram positive and Gram negative bacteria (*Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Salmonella enterica* and Cold and flu viruses (respiratory syncytial virus, Influenza Virus H1N1). Refer to Usage Table for the appropriate doses and contact times.

**Notice to User:** This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

### GENERAL DISINFECTION PERFORMANCE

**PUR TABS** is a general disinfectant effective against *Staphylococcus aureus*, *Salmonella enterica*, *Pseudomonas aeruginosa* and cold and flu respiratory syncytial virus, Influenza H1N1 when used at the dosage and contact time as detailed in the usage table.

### GENERAL DISINFECTION DIRECTIONS

Prepare a 1076 ppm solution. Apply to pre-cleaned surface with mop, cloth, sponge, brush, wipe, or coarse trigger sprayer. Allow surface to remain wet for 10 minutes. Allow to air dry.

### DISINFECTION/VIRUCIDAL<sup>†</sup> DIRECTIONS:

Prepare solution strength as required, refer to Usage Table for correct doses and contact times; refer to Dilution Chart for solution preparation. Apply use solution to pre-cleaned, hard, non-porous, inanimate surfaces with brush, spray device, sponge, cloth, or mop to wet all surfaces thoroughly. Allow to remain wet for contact time as indicated in the Usage Table, then remove product by wiping with brush, sponge, or cloth.

For sprayer applications using a spray device, spray at appropriate distance from surface depending on sprayer type (6 - 8 inches for spray bottles). Allow to remain wet for contact time as indicated in the Usage Table, then remove product by rubbing with brush, sponge, wipe or cloth or allow to air dry. Do not breathe spray mist.

Before using this product, food products and packaging materials must be removed from the room or carefully protected.

## KILLS HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1), HEPATITIS A VIRUS, AND HEPATITIS B VIRUS AND HEPATITIS C VIRUS ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS

PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS). Kills Human Immunodeficiency Virus Type 1 (HIV-1), Hepatitis A virus and Hepatitis B virus at 1076 ppm active chlorine solution in 10 minutes. Kills, Human Immunodeficiency Virus Type 1 (HIV-1), Hepatitis A virus, Hepatitis B virus and Hepatitis C virus at 4306 ppm active chlorine solution in 1 minute. Refer to Usage Table for correct doses and contact times. Refer to Dilution Chart for solution preparation.

## SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST Human Immunodeficiency Virus Type 1 (HIV-1) OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

**PERSONAL PROTECTION:** Specific barrier protection items to be used when handling items soiled with blood or body fluids are disposable latex gloves, gowns, masks, and eye coverings.

**CLEANING PROCEDURE:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of PUR TABS. This cleaning process may be accomplished with any cleaning solution including PUR TABS.

**DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids should be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.

**CONTACT TIME:** Leave surfaces wet for 10 minutes when using a 1076 ppm solution. Leave surfaces wet for 1 minute when using a 4306 ppm solution. Refer to Usage Table for correct doses and contact times. Refer to Dilution Chart for solution preparation

### ANIMAL PATHOGENS:

When used at dosage and contact times as outlined in the Usage Table, PUR TABS is effective against the following animal pathogens: Canine Parvovirus, Herpes simplex virus type 1<sup>†</sup> Newcastle Disease Virus, Pseudorabies, Feline Calicivirus, Norovirus, Canine Distemper virus, Infectious Canine hepatitis<sup>†</sup>, Teschen/Talfan disease<sup>†</sup>, Avian influenza Virus<sup>†</sup>, Porcine parvovirus<sup>†</sup>, Runting & Stunting virus (tenosynovitis)<sup>†</sup>, *Actinobacillus pleuropneumoniae*<sup>†</sup>, *Bordetella bronchiseptica* (rhinitis)<sup>†</sup>, *Brachyspira* (Treponema/Serpulina)<sup>†</sup>, Hyodysenteriae (swine dysentery)<sup>†</sup>, Gumboro disease<sup>†</sup>, Porcine Epidemic Diarrhea Virus<sup>†</sup>, *Streptococcus uberis*<sup>†</sup>, Transmissible gastroenteritis (TGE)<sup>†</sup>, Swine Vesicular disease<sup>†</sup>, African swine fever<sup>†</sup>, Hog cholera/Classical swine fever<sup>†</sup>, Avipox (fowl pox)<sup>†</sup>, Respiratory syncytial virus<sup>†</sup>, Bovine Viral Diarrhea Virus<sup>†</sup> and Porcine epidemic diarrhea virus<sup>†</sup>. Re-apply product as necessary to ensure surface remains wet.

<sup>†</sup> Note: This use has not been approved by the California DPR

## SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION IN ANIMAL HOUSING FACILITIES:

1. Remove all animals and feed from premises, vehicles, and enclosures.
2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with water.
5. Saturate all surfaces with appropriate solution strength for the appropriate contact time, refer to Usage Table for correct dose and contact time, and to Dilution Chart for solution preparation.
6. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
7. Ventilate buildings, cars, boats, and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set, or dried.
8. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers with soap or detergent, and allow to air dry before reuse.

**PUR TABS is also effective as a Healthcare disinfectant for critical areas potentially contaminated with C.diff spores when used at a level of 2153ppm available chlorine disinfectant solution. A 10 minute contact time is required.**

## DISINFECTION FOR SURFACES CONTAMINATED WITH CLOSTRIDIUM DIFFICILE

### Special Label Instructions for Cleaning Prior to Disinfection against Clostridium difficile spores:

**Personal Protection:** Wear appropriate barrier protection such as gloves, gowns, masks or eye covering.

**Cleaning Procedure:** Fecal matter/waste must be thoroughly cleaned from surfaces/objects before disinfection by application with a clean cloth, mop, and/or sponge saturated with the disinfectant product. Cleaning is to include vigorous wiping and/or scrubbing, until all visible soil is removed. Special attention is needed for high-touch surfaces. Surfaces in patient rooms are to be cleaned in an appropriate manner, such as from right to left or left to right on horizontal surfaces, and top to bottom on vertical surfaces, to minimize spreading of the spores. Restrooms are to be cleaned last. Do not reuse soiled cloths.

**Infectious Materials Disposal:** Materials used in the cleaning process that may contain feces/wastes are to be disposed of immediately in accordance with local regulations for infectious materials disposal.

**Directions for Use:** Prepare a 2153 ppm solution; refer to Dilution Chart. Apply to pre-cleaned surface with mop, cloth, sponge, brush, wipe, or mechanical sprayer. Allow surface to remain wet for 10 minutes. Allow to air dry.

### To Pre-clean Instruments Prior to Terminal Sterilization/High Level Disinfection

Prepare a 2153 ppm solution. As a pre-cleaning spray- Place instruments into a suitable container. Spray PUR TABS onto instruments to thoroughly wet all surfaces. Let stand for up to 10 minutes. Rinse instruments.

**As a pre-cleaning immersion solution –** Fill appropriate size container with a sufficient amount of PUR TABS to completely submerge instruments. Place instruments into the container of PUR TABS, cover, and allow to soak for up to 10 minutes. Remove and rinse and follow with an appropriate cleaning and disinfecting process. Change solution daily.

**As a manual instrument cleaner –** Thoroughly pre-rinse dirty instruments under running water to remove gross debris. Immerse pre-rinsed instruments into an appropriate size container filled with PUR TABS. Scrub instruments using a stiff bristle brush until visibly clean. Submerge instruments while scrubbing. Rinse instruments thoroughly. Change solution daily. Follow with an appropriate disinfection process. Cleaning of critical and semi-critical devices must be followed by an appropriate terminal sterilization/high level disinfection process.

**To Disinfect Non-Critical Pre-Cleaned Instruments –** Instruments must be thoroughly pre-cleaned to remove excess organic debris rinsed and rough dried (Clean and rinse lumens of hollow instruments before filling with PUR TABS or before immersion). Immersion method using a soaking tray: Immerse instruments into PUR TABS and let stand for ten or 10 minutes. Change solution for each use. Spray method – Spray all surfaces of instruments with PUR TABS until thoroughly wet. Let stand for 10 minutes.

## HEALTHCARE DISINFECTION WITHOUT PRECLEAN PERFORMANCE

PUR TABS is a Healthcare disinfectant when used at a level of 4,306 ppm available chlorine disinfectant solution. It is effective against Norovirus, Coxsackievirus and *Aspergillus fumigatus* with a 1 minute contact time. It is effective against *Salmonella enterica*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, carbapenem resistant *Klebsiella pneumoniae*, *Acinetobacter baumannii*, vancomycin resistant *Enterococcus faecalis*, *Staphylococcus aureus* - methicillin-resistant (MRSA), & glycopeptide-resistant (GRSA), *Streptococcus pneumoniae* and, *Candida albicans* with a 4 minute contact time.

## HEALTHCARE DISINFECTION WITHOUT PRECLEAN DIRECTIONS

Prepare a 4,306 ppm solution; refer to Dilution Chart. Apply to surface with mop, cloth, sponge, brush, wipe, or mechanical sprayer. Allow surface to remain wet for the appropriate contact time. Refer to Usage Table and HEALTHCARE DISINFECTION WITHOUT PRECLEAN PERFORMANCE.

PUR TABS is also effective as a Healthcare disinfectant for critical areas potentially contaminated with *Mycobacterium bovis* (Tb) when used at a level of 5382ppm available chlorine disinfectant solution. A 4 minute contact time is required.

## DISINFECTION FOR SURFACES CONTAMINATED WITH MYCOBACTERIUM BOVIS (Tb) IN 4 MINUTES at 20°C (68°F)

### Special Label Instructions for Cleaning Prior to Disinfection against Mycobacterium bovis (Tb):

This product when used as directed below is effective against *Mycobacterium bovis* (Tb). This product can be used on hard non-porous surfaces in commercial institutional hospital and other premises (including kitchens, bathrooms, nurseries, sick rooms, laundry rooms, eating establishments, pet kennels, and veterinary premises). To disinfect hard non-porous surfaces, first clean surface by removing gross filth (loose dirt, debris, food materials etc.). Prepare a 5,382 ppm available chlorine solution. Thoroughly wet surface with the solution and allow it to remain in contact with the surface for 4 minutes. Allow to air dry.

## SANITIZER FOR FOOD AND BEVERAGE PROCESSING AND FOOD HANDLING OPERATIONS

Prepare a 100 ppm solution; refer to dilution chart for the number of tablets to use.

This product is recommended for sanitizing all types of hard, non-porous equipment and utensils used

# PUR TABS™

## EFFERVESCENT SANITIZING / DISINFECTION TABLETS



in food processing and canning plants, bottling plants, breweries, fish processing plants, meat and poultry processing plants, milk handling and processing plants, stores, restaurant and institutional dining establishments. Use a 100 ppm available chlorine solution (refer to Dilution Chart) to sanitize previously cleaned processing and packaging equipment. Allow at least a one minute contact time before draining. Allow adequate draining before contact with beverages.

### SANITIZING HARD, NON-POROUS SURFACES, DISHES, GLASSES, FOOD PROCESSING EQUIPMENT AND UTENSILS, DAIRY AND BREWERY EQUIPMENT AND UTENSILS

**Prepare a 100 ppm solution; refer to dilution chart for the number of tablets to use.**  
This product is an effective sanitizing agent. Treatment with this product throughout food and beverage processing and food handling operations can help ensure the quality of the final product.

#### FOOD CONTACT SANITIZING DIRECTIONS

##### HANDWASHING OF ITEMS IN A 3 COMPARTMENT SINK

1. Remove all gross food particles and soil by a preflush or prescrape and, when necessary, presoak treatment. Wash surfaces or objects with a good detergent or compatible cleaner, followed by a potable water rinse before application of the sanitizing solution.
2. Prepare a 100 ppm available chlorine sanitizing solution (refer to Dilution Chart).
3. Place equipment, utensils, dishes, glasses, etc. in the solution or apply the use solution to surfaces using a cloth, sponge, or coarse sprayer.
4. Allow to stand for at least 1 minute, drain the excess solution from the surface, and allow to air dry.
5. Fresh sanitizing solution must be prepared at least daily or more often if the solution becomes diluted or soiled.

##### HANDWASHING OF ITEMS IN A 2 COMPARTMENT SINK

1. Remove all gross food particles and soil by a preflush or prescrape and, when necessary, presoak treatment. Wash surfaces or objects with a good detergent or compatible cleaner,
2. Prepare a 100 ppm available chlorine sanitizing solution (refer to Dilution Chart).
3. Place equipment, utensils, dishes, glasses, etc. in the solution or apply the use solution to surfaces using a cloth, sponge, or coarse sprayer.
4. Allow to stand for at least 1 minute, drain the excess solution from the surface, and allow to air dry.
5. Fresh sanitizing solution must be prepared at least daily or more often if the solution becomes diluted or soiled.

##### MACHINE WASHING OF ITEMS

1. Remove all gross food particles and soil by a preflush or prescrape and, when necessary, presoak treatment. Wash surfaces or objects with a good detergent or compatible cleaner, followed by a potable water rinse before application of the sanitizing solution.
2. Prepare a 100 ppm available chlorine solution (refer to Dilution Chart).
3. Add the solution to the feed tank of immersion or spray type machines that can provide at least 1 minute contact time for sanitizing dishes, glasses, food processing equipment, or utensils. Allow to drain and air dry before use.
4. Promptly use the sanitizing solution after preparation. Discard unused solutions

The following directions for Use are not allowed in the state of California:

##### SHOE AND BOOT BATH DEODORIZER

To deodorize footwear worn in animal areas and in packaging and storage areas of food plants. Shoe and Boot baths containing one inch of freshly made 100 ppm available chlorine solution (refer to Dilution Chart) should be placed at all entrances to buildings, hatcheries, and at all the entrances to the production and packaging rooms. Scrape waterproof shoes and boots and place into solution for at least 1 minute prior to entering area. Change the solution in the bath at least daily or sooner if solution appears diluted or dirty.

##### MILK HANDLING AND PROCESSING EQUIPMENT

This product can be used on dairy farms and in plants processing milk, cream, ice cream, and cheese. Rinse milking machines, utensils, and all equipment with cold water to remove excess milk. Clean with a suitable detergent cleaning product and or water as appropriate] and rinse prior to sanitizing. To sanitize, spray or rinse all pre-cleaned surfaces with 100 ppm available chlorine solution (refer to Dilution Chart). Allow at least a 1 minute contact time before draining. Allow adequate draining before contact with dairy products. It is important to clean out large deposits of milk or other organic matter before sanitizing. A sharp decline in the available chlorine content of the sanitizer following circulation through milk processing equipment is usually regarded as evidence of inadequate cleaning of the equipment and should be promptly investigated.

The following directions for Use are not allowed in the state of California:

##### SANITIZING APPLICATION METHODS

Prepare a 100 ppm solution; refer to Dilution Chart for the number of tablets to use. Freshly prepare all sanitizing solutions. Test solutions during use to ensure the concentration does not drop below the recommended level. Keep in properly labeled containers to protect against contamination. Discard unused solutions.

##### GENERAL RINSE METHOD

Prepare a solutions containing 100 ppm available chlorine (refer to Dilution Chart) sanitize plant floors, walls and ceilings and also control odors in refrigerated areas and drain platforms. Generously flush or swab surfaces with the solution. After one minute contact time allow solution to drain and then air dry.

##### DISINFECTION OF DRINKING WATER IN EMERGENCY/PUBLIC/INDIVIDUAL SYSTEMS EMERGENCY DRINKING WATER

Use PUR TABS to disinfect raw or pre-treated (settled, coagulated, and/or filtered) human and domestic animal drinking supplies on an emergency basis as defined in 40 CFR, Part 165-179. The treated water source may be a river, lake, well, cistern or similar system. The treated water should be clear and free of dirt and organic debris to obtain the optimum disinfection results. If the water source is cloudy and contains dirt and organic debris, the water should be in holding tanks or pond, treated with coagulating agents and filtered to remove dirt and organic debris.

Refer to dilution chart for the number of tablets to use to achieve available chlorine concentration of 10 ppm. Allow water to stand for seven to fifteen minutes before use. Maintain 1 ppm available chlorine residual, as determined by a reliable chlorine test kit, to ensure disinfection.

##### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in a cool, dry, well-ventilated area at a temperature below 40°C/104°F. Avoid moisture getting into container.

**PESTICIDE DISPOSAL:** Pesticides may be acutely hazardous. Wastes resulting from the use of this product must be disposed of on-site, or at an approved waste disposal facility.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container.

Offer for recycling if available or puncture and dispose of in a sanitary landfill.

Sold by:



EarthSafe Chemical Alternatives  
Braintree, MA 02184  
1-866-666-2305  
www.evaclean.com

EPA Reg. No. 71847-6-91524  
EPA Est. No. 71847-IRL-001

#### USAGE TABLE:

PATHOGEN	MINIMUM DOSE REQUIRED (PPM)	MINIMUM CONTACT TIME REQUIRED (MINUTES)
<b>Food Contact Sanitizer Claims</b>		
<i>Staphylococcus aureus</i> (ATCC 6538)	100 ppm	1 minute
<i>Salmonella enterica</i> (ATCC 6539)	100 ppm	1 minute
<b>Disinfection Claims - Bacteria</b>		
<i>Staphylococcus aureus</i> (ATCC 6538)	a) 1076 ppm b) 4306 ppm	a) 10 minutes b) 4 minutes
<i>Staphylococcus aureus</i> - methicillin resistant (MRSA) & glycopeptide-resistant (GRSA) (ATCC 33592)	a) 1076 ppm b) 4306 ppm	a) 10 minutes b) 4 minutes
<i>Staphylococcus epidermidis</i> (ATCC 51624)	1076 ppm	10 minutes
<i>Salmonella enterica</i> (ATCC 10708)	a) 1076 ppm b) 4306 ppm	a) 10 minutes b) 4 minutes
<i>Pseudomonas aeruginosa</i> (ATCC 15442)	a) 1076 ppm b) 4306 ppm	a) 10 minutes b) 4 minutes
<i>Streptococcus pneumoniae</i> (ATCC 6305)	4306 ppm	4 minutes
<i>Escherichia coli</i> O157:H7 (ATCC 35150)	1076 ppm	10 minutes
<i>Acinetobacter baumannii</i> (ATCC BAA-1709)	4306 ppm	4 minutes
Vancomycin resistant <i>Enterococcus faecalis</i> (ATCC 51575)	a) 1076 ppm b) 4306 ppm	a) 10 minutes b) 4 minutes
Carbapenem resistant <i>Klebsiella pneumoniae</i> (ATCC BAA-1705)	4306 ppm	4 minutes
<i>Klebsiella pneumoniae</i> (ATCC 4352)	1076 ppm	10 minutes
<b>Virucidal Claims</b>		
Respiratory syncytial virus <sup>†</sup> (ATCC VR-26)	538 ppm	10 minutes
Rhinovirus Type 14 <sup>†</sup> (ATCC VR-284)	1076 ppm	10 minutes
Influenza Virus H1N1 <sup>†</sup> (ATCC VR-99)	1076 ppm	10 minutes
Human Immunodeficiency Virus Type 1 (HIV-1) <sup>†</sup> (Strain IIIB)	a) 1076 ppm b) 4306 ppm	a) 10 minutes b) 1 minute
Hepatitis A virus <sup>†</sup> (Strain HM175/18f)	a) 1076 ppm b) 4306 ppm	a) 10 minutes b) 1 minute
Hepatitis B virus <sup>†</sup> (Duck Hepatitis B (DHBV))	a) 1076 ppm b) 4306 ppm	a) 10 minutes b) 1 minute
Hepatitis C virus <sup>†</sup> (Bovine Viral Diarrhea Virus Strain NADL - surrogate for Hepatitis C virus)	4306 ppm	1 minute
Avian influenza A (H5N1) <sup>†</sup> (CDC #2006719965)	4306 ppm	1 minute
Norovirus <sup>†</sup> (ATCC VR-782)	2153 ppm	1 minute
Poliovirus Type 1 <sup>†</sup> (ATCC VR-1000)	1076 ppm	10 minutes
Coxsackievirus B3 <sup>†</sup> (ATCC VR- 30)	4306 ppm	1 minute
Herpes simplex virus type 1 <sup>†</sup> (ATCC VR-733)	1076 ppm	10 minutes
<b>Fungicidal/Yeasticidal Claims</b>		
<i>Aspergillus fumigatus</i> (ATCC 36607)	4306 ppm	1 minute
<i>Candida albicans</i> (ATCC 10231)	4306 ppm	4 minutes
<i>Trichophyton interdigitale</i> (ATCC 9533)	1076 ppm	10 minutes
<b>Clostridium difficile Claims</b>		
<i>Clostridium difficile</i> spores (ATCC 43598)	a) 2153 ppm b) 4306 ppm	a) 10 minutes b) 4 minutes
<b>Mycobactericidal Claims</b>		
<i>Mycobacterium bovis</i> (TB) (ATCC 35743)	5382 ppm	4 minutes
<b>Animal Pathogens<sup>†</sup></b>		
Canine Parvovirus <sup>†</sup> (ATCC VR-2017)	1076 ppm	10 minutes
Herpes simplex virus type 1 <sup>*†</sup> (ATCC VR-733)	1076 ppm	10 minutes
Newcastle Disease Virus <sup>†</sup> (ATCC VR-180)	1076 ppm	10 minutes
Pseudorabies <sup>†</sup> (ATCC VR-135)	1076 ppm	10 minutes
Feline Calicivirus <sup>†</sup> (ATCC VR-782)	1076 ppm 2153 ppm	10 minutes 1 minute
Canine Distemper virus <sup>†</sup> (ATCC VR-128)	1076 ppm	10 minutes
Infectious Canine hepatitis <sup>*†</sup> (ATCC VR 293)	1076 ppm	10 minutes
Teschen/Talfan disease <sup>*†</sup> (ATCC VR-669)	1076 ppm	10 minutes
Influenza Virus H1N1 <sup>†</sup> (ATCC VR-99)	1076 ppm	10 minutes
Avian influenza virus H5N1 <sup>*†</sup> (ATCC VR-1608)	4306 ppm	1 minute
Porcine parvovirus <sup>*†</sup> (ATCC VR-742)	1076 ppm	10 minutes
Runting & Stunting virus (tenosynovitis) <sup>*†</sup> (ATCC VR- 2449) (ATCC VR-21)	1076 ppm	10 minutes
<i>Actinobacillus pleuropneumoniae</i> <sup>*†</sup> (NCTC 12370) (ATCC 27088)	1076 ppm	10 minutes
<i>Bordetella bronchiseptica</i> (rhinitis) <sup>*†</sup> (ATCC 19)	1076 ppm	10 minutes
Brachyspira (Treponema/Serpulina) <sup>*†</sup> (ATCC 27164)	1076 ppm	10 minutes
Hyodysenteriae (swine dysentery) <sup>*†</sup> (ATCC 27164)	1076 ppm	10 minutes
Gumboro disease <sup>*†</sup> (ATCC VR-478)	1076 ppm	10 minutes
<i>Streptococcus uberis</i> <sup>*†</sup> (ATCC 9927)	1076 ppm	10 minutes
Transmissible gastroenteritis (TGE) <sup>*†</sup> (ATCC VR-743)	1076 ppm	30 minutes
Swine Vesicular disease <sup>*†</sup> (ATCC VR-158)	1076 ppm	30 minutes
African swine fever <sup>*†</sup> (ASFV)	1076 ppm	30 minutes
Hog cholera/Classical swine fever <sup>*†</sup> (CSFV)	1076 ppm	30 minutes
Avipox (fowl pox) <sup>*†</sup> (FPV)	1076 ppm	30 minutes
Respiratory syncytial virus <sup>*†</sup> (ATCC VR-26)	538 ppm	10 minutes
Bovine Viral Diarrhea Virus <sup>*†</sup> (Strain NADL)	4306 ppm	1 minute
Porcine epidemic diarrhea virus <sup>*†</sup> (Strain Colorado)	1076 ppm	10 minutes

<sup>†</sup>Note: This use has not been approved by the California DPR

<sup>\*</sup>Note: these organisms not approved by the state of California